

# ON THE EXPONENTIAL DIOPHANTINE EQUATION

VÎRGOLICI, Horia

Faculty of Mathematics and Informatics

Spiru Haret University

hvirgolici@yahoo.com

## Abstract

*The aim of this paper is to give all nonnegative solutions  $(x, y, z)$  to the equation  $2^x + p^y = 1009^z$ , where  $p$  is a positive rational prime number with  $3 \leq p \leq 997$  (167 equations)*

**Keywords:** *exponential Diophantine equation, Lebesgue-Nagell equation, Catalan equation*

**AMS Classification:** 11D61, 11Y50

## 1. Introduction

It is known that the equation

$$a^x + b^y = c^z \tag{1}$$

where  $a, b, c$  are prime numbers, has only finitely many solutions, but there is no algorithm to compute all the solutions  $(x, y, z)$ . Some particular cases were treated: Nagell [8] found all solutions for  $\max(a, b, c) = 7$ , Makowski [5], Hadano [4], Uchiyama [12], Qi Sun and Xiaoming Zhou [10] and Xiaozhuo Yang [13] determined all solutions for  $11 \leq \max(a, b, c) \leq 23$ . Cao [3] found all solutions for  $29 \leq \max(a, b, c) \leq 97$  (60 solutions in total).

The aim of this paper is to give all nonnegative solutions to equation (1) for  $a = 2, c = 1009$  (1009 representing the first prime number  $> 1000$ ) and  $b$  is a rational prime number,  $3 \leq b \leq 997$ . The main results are given by the following

**Theorem 1.** *The only equations  $2^x + p^y = 1009^z$ , with  $p$  rational prime,  $3 \leq p \leq 1009$ , which admit nonnegative solutions  $(x, y, z)$ , are:*

- $2^x + 881^y = 1009^z$ , which has the solution  $(x, y, z) = (7, 1, 1)$ .
- $2^x + 977^y = 1009^z$ , which has the solution  $(x, y, z) = (5, 1, 1)$ .

The proof for Theorem 1 is given in subsection 3.

## 2. Preliminaries

Below we present some theorems which establish the maximum number of solutions for the equation (1). Theorem 2 shows that the Catalan's equation has only one solution and gives this solution.

**Theorem 2.** ([1, 6, 7]) *Equation (named Catalan's equation)*

$$a^x - b^y = 1 \tag{2}$$

*has no solutions in integers  $a, b, x, y > 1$  other than  $3^2 - 2^3 = 1$ .*

Theorem 3 is concerned with equation  $a^x + b^y = p^z$ ,  $p$  rational prime, and proves that this equation has at most two solutions, except some particular equations, which have three solutions.

**Theorem 3.** ([9, Lemma 6, p.163]) *The equation*

$$a^x + b^y = p^z \tag{3}$$

*has at most one solution when the parity of  $x$  and  $y$  are preassigned, except for three choices of  $(a, b, p)$  taking  $(a < b)$  :  $(3, 5, 2)$ ,  $(3, 13, 2)$ ,  $(3, 10, 13)$ .*

## 3. Proofs of the main results

We give the proof for Theorem 1, which treats the equation

$$2^x + p^y = 1009^z \tag{4}$$

If  $y = 0$ , the equation (4) has no solutions, due to Theorem 2. So we consider  $y > 0$ . Many of the equations (4) have no solutions  $(x, y, z)$  due to:

**Lemma 1.** *If  $p \equiv 1, 5, 7, 11, 13, 19, 23, 25, 29, 31, 35, 37, 43 \pmod{48}$  or  $p \equiv 1 \pmod{7}$ , then the equation (4) has no nonnegative solutions  $(x, y, z)$ .*

*Proof:* It is obvious, taking the equation (4) mod 48 and mod 7, respectively.

Thus, we discuss the following equations, in which  $p$  doesn't comply with the conditions in Lemma 1:

- i/  $2^x + p^y = 1009^z$ , where  $p \in \{89, 137, 233, 521, 569, 761, 809, 857\}$ ; by taking the equation mod 48, it results that  $x = 3$ ; by taking the equation mod 7, it results that there are no solutions.
- ii/  $2^x + p^y = 1009^z$ , where  $p \in \{191, 431, 863\}$ ; by taking the equation mod 48, it results that  $x = 1$ ; by taking the equation mod 7, it results that there are no solutions.
- iii/  $2^x + 3^y = 1009^z$ , which has no solutions by taking the equation mod 60.
- iv/  $2^x + 17^y = 1009^z$ , which has no solutions by taking the equation mod 3, which leads to  $x$  and  $y$  odd, and mod 17, which becomes impossible with  $x$  and  $y$  odd.

- v/  $2^x + 41^y = 1009^z$ ; by taking the equation mod 48, it results that  $x=3$ , so the equation becomes  $8 + 41^y = 1009^z$ , which has no solutions by taking the equation mod 7.
- vi/  $2^x + 47^y = 1009^z$ ; by taking the equation mod 48, it results that  $x=1$ , so the equation becomes  $2 + 47^y = 1009^z$ , which has no solutions, by taking the equation mod 3 and mod 13.
- vii/  $2^x + 257^y = 1009^z$ ; by taking the equation mod 3, it results that  $x$  and  $y$  are odd; by taking the equation mod 32, one gets  $x \geq 5$  and  $z$  is even; by taking the equation mod 5, it results that  $x \equiv 3 \pmod{4}$  and  $y \equiv 3 \pmod{4}$ ; by taking the equation mod 13, it results that  $x \equiv 3 \pmod{12}$ ,  $y \equiv 11 \pmod{12}$  and  $z \equiv 2 \pmod{4}$ ; by taking the equation mod 7, one concludes that there are no solutions in this case.
- viii/  $2^x + 353^y = 1009^z$ ; by taking the equation mod 3, it results that  $x$  and  $y$  are odd; by taking the equation mod 7, it results that  $x \equiv 1 \pmod{3}$  and  $y \equiv 3 \pmod{6}$ ; by taking the equation mod 9 and taking into account the relations above, it results that  $x \equiv 1 \pmod{6}$ ; by taking mod 13 and taking into account the relations obtained till now, the equation has no solutions.
- ix/  $2^x + 383^y = 1009^z$ ; by taking the equation mod 24, it results that  $x = 1$  and  $y$  is odd, so the equation becomes  $2 + 383^y = 1009^z$ ; by taking this equation mod 5, it results that  $y \equiv 3 \pmod{4}$  and  $z$  is odd; by taking the equation mod 13, one concludes that there are no solutions in this case.
- x/  $2^x + 401^y = 1009^z$ ; by taking the equation mod 7, it results that  $x \equiv 2 \pmod{3}$  and  $y \equiv 2 \pmod{3}$ ; by taking the equation mod 9 and taking into account the relations above, it results that the equation has no solutions.
- xi/  $2^x + 479^y = 1009^z$ ; by taking the equation mod 48, it results that  $x=1$ ; so the equation becomes  $2 + 479^y = 1009^z$ ; by taking the equation mod 9, it results that  $y \equiv 3 \pmod{6}$ ; taking this relation into consideration and taking the equation mod 13, it results that the equation has no solutions.
- xii/  $2^x + 593^y = 1009^z$ ; by taking the equation mod 3, it results that  $x$  and  $y$  are odd; by taking the equation mod 9, it results that  $x \equiv 1 \pmod{6}$ ; taking the equation mod 13, it results that the equation has no solutions.
- xiii/  $2^x + 641^y = 1009^z$ ; by taking the equation mod 32, it results that one has either  $x = 4$  and  $z$  odd or  $x \geq 5$  and  $z$  even; in both cases, by taking the equation mod 5, it results that the equation has no solutions.
- xiv/  $2^x + 719^y = 1009^z$ ; by taking the equation mod 48, it results that  $x=1$ , so the equation becomes  $2 + 719^y = 1009^z$ ; by taking this equation mod 1009, it results that the equation has no solutions.
- xv/  $2^x + 881^y = 1009^z$ ; by taking the equation mod 3, it results that  $x$  and  $y$  are odd; the equation has the solution  $(x,y,z)=(7,1,1)$  and it is the only one, due to Theorem 3.

- xvi/  $2^x + 929^y = 1009^z$ ; by taking the equation mod 3, it results that  $x$  and  $y$  are odd; by taking the equation mod 7, it results that  $x \equiv 1 \pmod{6}$  and  $y \equiv 3 \pmod{6}$ ; taking now the equation mod 13, it results that it has no solutions.
- xvii/  $2^x + 977^y = 1009^z$ ; by taking the equation mod 3, it results that  $x$  and  $y$  are odd; the equation has the solution  $(x,y,z)=(5,1,1)$  and it is the only solution, due to Theorem 3.

## References

1. Bilu, Yu.F., *Catalan's Conjecture (after Mihăilescu)*, "Séminaire Bourbaki, Exposé" 909, 55ème année, 2002-2003.
2. Bugeaud, Y., Mignotte, M., Siksek, S., *Classical And Modular Approaches To Exponential Diophantine Equations. II. The Lebesgue - Nagell Equation*, "Compos. Math.", 142, 31-62, 2006.
3. Cao, Z. F., *On the Diophantine Equation  $a^x = b^y + c^z$* , "Chinese Sci. Bull.", I, 32, 1519-1521, 1987; II, ibid. 33, 237 (in Chinese), 1988.
4. Hadano, T., *On the Diophantine Equation  $a^x = b^y + c^z$* , "Math. J. Okayama Univ.", 19, 25-29, 1976/77.
5. Makowski, A., *On the Diophantine Equation  $2^x + 11^y = 5^z$* , "Nord. Mat. Tidskr.", 7, 81-96, 1959.
6. Mihăilescu, P., *A Class Number Free Criterion for Catalan's Conjecture*, "J. of Number Theory", 99, 225-231, 2003.
7. Mihăilescu, P., *Primary Cyclotomic Units and a Proof of Catalan's Conjecture*, "Journal für die reine und angewandte Mathematik", 572, 167-195, 2004.
8. Nagell, T., *Sur une classe d'équations exponentielles*, "Ark. Mat.", 3, 569-582, 1958.
9. Scott, R., *On the Equations  $p^x - b^y = c$  and  $a^x + b^y = c^z$* , "J. of Number Theory", 44, 153-165, 1993.
10. Sun, Q., Zhou, X. M., *On the Diophantine Equation  $a^x = b^y + c^z$* , "Chinese Sci. Bull.", 29, 61, 1984 (in Chinese).
11. Terai, N., *Applications of a Lower Bound for Linear Forms in two Logarithms to Exponential Diophantine Equations*, "Acta Arithmetica XC", 1, 17-35, 1999.
12. Uchiyama, S., *On the Diophantine Equation  $2^x = 3^y + 13^z$* , "Math. J. Okayama Univ.", 19, 31-38, 1976/77.
13. Yang, X. Z., *On the Diophantine Equation  $a^x = b^y + c^z$* , "Sichuan Daxue Xuebao", 4, 151-158, 1985 (in Chinese).